







- Excellent productivity.
- Medium early variety.
- Big oval to long oval tubers.
- White flesh.

## **PLANT & TUBER**

## **PESTS & DISEASES**

FOLIAGE MATURITY	MEDIUM EARLY	RESISTANCE TO FOLIAGE LATE BLIGHT	FAIRLY LOW
EMERGENCE	MODERATE	RESISTANCE TO TUBER LATE BLIGHT	FAIRLY LOW
PLANT VIGOUR	HIGH	RESISTANCE TO PITTED COMMON SCAB	MODERATE
SKIN COLOUR	PALE YELLOW	RESISTANCE TO NETTED COMMON SCAB	MODERATE
FLESH COLOUR	WHITE	RESISTANCE TO PVY	FAIRLY LOW
TUBER SHAPE	OVAL TO LONG OVAL	RESISTANCE TO PCN G.rostochiensis (Ro1-4)	LOW
SHAPE UNIFORMITY	GOOD	RESISTANCE TO BLACK WART	HIGH
EYE DEPTH	SHALLOW		
TUBER SIZE	LARGE		

## **QUALITY & USE**

YIELD
DRY MATTER CONTENT
COOKING TYPE
DISINTEGRATION
DARKENING AFTER COOKING
FRY COLOUR at harvest
STORAGE ABILITY

VERY HIGH LOW A VERY SLIGHT NONE MEDIUM FAIRLY HIGH

## PHYSIOLOGY

DORMANCY	MEDIUM
RESISTANCE TO EXTERNAL DAMAGE	FAILRY HIGH
RESISTANCE TO INTERNAL DAMAGE	FAIRLY HIGH
RESISTANCE TO INTERNAL RUST SPOT	FAIRLY HIGH
RESISTANCE TO DESPROUTING	HIGH

The information provided in this document is only indicative and may vary according to growing and climatic conditions but also agronomic techniques. Updated : 2025-06-25